Risk Factors for Cognitive Decline: Delaware

Based on population-level evidence, the six health conditions and behaviors included here increase risk for cognitive decline — and may also increase risk of dementia.

UBLIC HEALT CENTER OF EXCELLENCE

DEMENTIA RISK

Data are from the Behavioral Risk Factor Surveillance System (BRFSS).

Percent With at Least One of Five Risk Factors (excluding sleep)

## 40.1% 41.7% Midlife Hypertension Physical Inactivity (age 45-64) 40.9% 13.3% Midlife Obesity Diabetes (age 45-64) 12.6% 11.8% Smoking Poor Sleep (age 45 and older) (<6 hours/night)

Prevalence of Six Risk Factors

## Percent With Any of Five Risk Factors (excluding sleep)

None 28.6%	One 33.7%	Two or more 37.7%

The Public Health Center of Excellence on Dementia Risk Reduction is made possible by the Centers for Disease Control and Prevention (CDC) of the U.S. Department of Health and Human Services (HHS) as part of a financial assistance award totaling \$3,707,435. The contents of this fact sheet are those of the Alzheimer's Association and do not necessarily represent the official views of, nor an endorsement by, CDC, HHS, or the U.S. government.

Updated: December 2024

CenterOfExcellence@alz.org

## alz.org/publichealth